



Biological Monitoring

Quarterly Compliance Calendar

How often should I perform biological indicator monitoring (spore testing)?

The CDC recommends testing **at least weekly** with a biological monitor for each sterilizer to ensure proper functioning. Users should follow the manufacturer's directions concerning the appropriate placement of the BI in the sterilizer. In addition to conducting routine biological monitoring, equipment users should perform biological monitoring:

- Whenever a new type of packaging material or tray is used.
- After training new sterilization personnel.
- After any sterilizer repair or maintenance.
- After any change in the sterilizer loading procedures.

Circle the date each week that BI testing was performed

JANUARY

S	M	T	W	T	F	S
			1	2	3	4
5	6	7	8	9	10	11
12	13	14	15	16	17	18
19	20	21	22	23	24	25
26	27	28	29	30	31	

FEBRUARY

S	M	T	W	T	F	S
						1
2	3	4	5	6	7	8
9	10	11	12	13	14	15
16	17	18	19	20	21	22
23	24	25	26	27	28	

MARCH

S	M	T	W	T	F	S
						1
2	3	4	5	6	7	8
9	10	11	12	13	14	15
16	17	18	19	20	21	22
23	24	25	26	27	28	29
30	31					

[illegible]

FOR STERILITY ASSURANCE:

EVERY LOAD: Sure-Check™ Sterilization Pouches

EVERY DAY: SteamPlus™ Integrating Indicators

EVERY WEEK: ConFirm™ Biological Monitoring Services

For test status or questions call customer care at: 1-800-722-1529 | CrosstexBMS.com



Hu-Friedy Mfg. Co., LLC, 1666 E. Touhy Ave., Des Plaines, IL 60018 | HuFriedyGroup.com
All company and product names are trademarks of Hu-Friedy Mfg. Co., LLC its affiliates or related companies, unless otherwise noted.

©2026 Hu-Friedy Mfg. Co., LLC. All rights reserved. 2411



HuFriedyGroup
The Best In Practice



Biological Monitoring

Quarterly Compliance Calendar

How often should I perform biological indicator monitoring (spore testing)?

The CDC recommends testing **at least weekly** with a biological monitor for each sterilizer to ensure proper functioning. Users should follow the manufacturer's directions concerning the appropriate placement of the BI in the sterilizer. In addition to conducting routine biological monitoring, equipment users should perform biological monitoring:

- Whenever a new type of packaging material or tray is used.
- After training new sterilization personnel.
- After any sterilizer repair or maintenance.
- After any change in the sterilizer loading procedures.

Circle the date each week that BI testing was performed

APRIL

S	M	T	W	T	F	S
		1	2	3	4	5
6	7	8	9	10	11	12
13	14	15	16	17	18	19
20	21	22	23	24	25	26
27	28	29	30			

MAY

S	M	T	W	T	F	S
				1	2	3
4	5	6	7	8	9	10
11	12	13	14	15	16	17
18	19	20	21	22	23	24
25	26	27	28	29	30	31

JUNE

S	M	T	W	T	F	S
1	2	3	4	5	6	7
8	9	10	11	12	13	14
15	16	17	18	19	20	21
22	23	24	25	26	27	28
29	30					

[illegible]

FOR STERILITY ASSURANCE:

EVERY LOAD: Sure-Check™ Sterilization Pouches

EVERY DAY: SteamPlus™ Integrating Indicators

EVERY WEEK: ConFirm™ Biological Monitoring Services

For test status or questions call customer care at: 1-800-722-1529 | CrosstexBMS.com



Hu-Friedy Mfg. Co., LLC, 1666 E. Touhy Ave., Des Plaines, IL 60018 | HuFriedyGroup.com
All company and product names are trademarks of Hu-Friedy Mfg. Co., LLC its affiliates or related companies, unless otherwise noted.

©2026 Hu-Friedy Mfg. Co., LLC. All rights reserved. 2411



HuFriedyGroup
The Best In Practice



Biological Monitoring

Quarterly Compliance Calendar

How often should I perform biological indicator monitoring (spore testing)?

The CDC recommends testing **at least weekly** with a biological monitor for each sterilizer to ensure proper functioning. Users should follow the manufacturer's directions concerning the appropriate placement of the BI in the sterilizer. In addition to conducting routine biological monitoring, equipment users should perform biological monitoring:

- Whenever a new type of packaging material or tray is used.
- After training new sterilization personnel.
- After any sterilizer repair or maintenance.
- After any change in the sterilizer loading procedures.

Circle the date each week that BI testing was performed

JULY

S	M	T	W	T	F	S
		1	2	3	4	5
6	7	8	9	10	11	12
13	14	15	16	17	18	19
20	21	22	23	24	25	26
27	28	29	30	31		

AUGUST

S	M	T	W	T	F	S
					1	2
3	4	5	6	7	8	9
10	11	12	13	14	15	16
17	18	19	20	21	22	23
24	25	26	27	28	29	30
31						

SEPTEMBER

S	M	T	W	T	F	S
	1	2	3	4	5	6
7	8	9	10	11	12	13
14	15	16	17	18	19	20
21	22	23	24	25	26	27
28	29	30				

[illegible]

FOR STERILITY ASSURANCE:

EVERY LOAD: Sure-Check™ Sterilization Pouches

EVERY DAY: SteamPlus™ Integrating Indicators

EVERY WEEK: ConForm™ Biological Monitoring Services

For test status or questions call customer care at: 1-800-722-1529 | CrosstexBMS.com



Hu-Friedy Mfg. Co., LLC, 1666 E. Touhy Ave., Des Plaines, IL 60018 | HuFriedyGroup.com
All company and product names are trademarks of Hu-Friedy Mfg. Co., LLC its affiliates or related companies, unless otherwise noted.

©2026 Hu-Friedy Mfg. Co., LLC. All rights reserved. 2411



HuFriedyGroup
The Best In Practice

A black circle with the year "2025" in white, bold, sans-serif font.

Biological Monitoring

Quarterly Compliance Calendar

How often should I perform biological indicator monitoring (spore testing)?

The CDC recommends testing **at least weekly** with a biological monitor for each sterilizer to ensure proper functioning. Users should follow the manufacturer's directions concerning the appropriate placement of the BI in the sterilizer. In addition to conducting routine biological monitoring, equipment users should perform biological monitoring:

- Whenever a new type of packaging material or tray is used.
- After training new sterilization personnel.
- After any sterilizer repair or maintenance.
- After any change in the sterilizer loading procedures.

Circle the date each week that BI testing was performed

OCTOBER

S	M	T	W	T	F	S
			1	2	3	4
5	6	7	8	9	10	11
12	13	14	15	16	17	18
19	20	21	22	23	24	25
26	27	28	29	30	31	

NOVEMBER

S	M	T	W	T	F	S
						1
2	3	4	5	6	7	8
9	10	11	12	13	14	15
16	17	18	19	20	21	22
23	24	25	26	27	28	29
30						

DECEMBER

S	M	T	W	T	F	S
	1	2	3	4	5	6
7	8	9	10	11	12	13
14	15	16	17	18	19	20
21	22	23	24	25	26	27
28	29	30	31			

[illegible]

FOR STERILITY ASSURANCE:

EVERY LOAD: Sure-Check™ Sterilization Pouches

EVERY DAY: SteamPlus™ Integrating Indicators

EVERY WEEK: ConFirm™ Biological Monitoring Services

For test status or questions call customer care at: 1-800-722-1529 | CrosstexBMS.com



Hu-Friedy Mfg. Co., LLC, 1666 E. Touhy Ave., Des Plaines, IL 60018 | HuFriedyGroup.com
All company and product names are trademarks of Hu-Friedy Mfg. Co., LLC its affiliates or related companies, unless otherwise noted.

©2026 Hu-Friedy Mfg. Co., LLC. All rights reserved. 2411



HuFriedyGroup
The Best In Practice